arXiv:hep-ph/9907202v2 17 Feb 2000

Inclusive Decays of Bottom Hadrons in New Formulation of Heavy Quark Effective Field Theory

Y.A Yan, Y.L Wu and W.Y Wang
Institute of Theoretical Physics, Chinese Academy of Sciences, Beijing 100080, China
(ylwu@itp.ac.cn)

Abstract

We apply the new formulation of heavy quark effective field theory (HQEFT) to the inclusive decays of bottom hadrons. The long-term ambiguity of using heavy quark mass or heavy hadron mass for inclusive decays is clarified within the framework of the new formulation of HQEFT. The $1/m_b$ order corrections are absent and contributions from $1/m_b^2$ terms are calculated in detail. This enables us to reliably extract the important CKM matrix element $|V_{cb}|$ from the inclusive semileptonic decay rates. The resulting lifetime ratios $\tau(B_s^0)/\tau(B^0)$ and $\tau(\Lambda_b)/\tau(B^0)$ are found to well agree with the experimental data. We also calculate in detail the inclusive semileptonic branching ratios and the ratios of the τ and β decay rates as well as the charm countings in the B^0, B_s^0 and Λ_b systems. For B^0 decays, all the observables are found to be consistent with the experimental data. More precise data for the B^0 decays and further experimental measurements for the B_s^0 and Λ_b systems will be very useful for testing the framework of new formulation of HQEFT at the level of higher order corrections.

PACS numbers: 12.39.Hg, 12.15.Hh, 13.20.He, 13.30.-a

Typeset using REVTEX

I. INTRODUCTION

In our previous paper [1], we have provided a more detailed study on a new formulation of HQEFT [2] and applied it to evaluate the weak transition matrix elements between the heavy hadrons containing a single heavy quark. Consequently, the new formulation of HQEFT has exhibited interesting features, such as: The Luke's theorem comes out automatically without imposing the equation of motion $iv\cdot DQ_v^+ = 0$; the form factors at zero recoil are found to be related to the meson masses, so that the most important relevant form factors at zero recoil can be fitted from the ground state meson masses; the number of universal form factors up to the order of $1/m_Q^2$ are less than the one in the usual HQET.

It is of interest to apply the new formulation of HQEFT [2] to the inclusive decays of bottom hadrons. The inclusive decays of bottom hadrons have been investigated in the recent years by several groups [3] − [10]. While it is well known that in the usual HQET there are still some problems which are not yet well understood. These problems mainly involve the following issues:

Firstly, the world average values for the lifetime ratios of bottom hadrons are [11]

$$\frac{\tau(B^-)}{\tau(B^0)} = 1.07 \pm 0.03, \tag{1.1a}$$

$$\frac{\tau(B_s^0)}{\tau(B^0)} = 0.94 \pm 0.04, \tag{1.1b}$$

$$\frac{\tau(\Lambda_b)}{\tau(B^0)} = 0.79 \pm 0.05. \tag{1.1c}$$

While the usual HQET prediction leads to a uniform lifetime for all the bottom hadrons when the nonspectator effects are neglected. The lifetime differences emerge from $1/m_Q^2$ order which is found to be small. One may expect that the $1/m_Q^3$ terms become dominant and provide about 20% contribution to the lifetime ratio $\tau(\Lambda_b)/\tau(B^0) = 0.79 \pm 0.05$ (or at least 10% corrections to their total decay rates). If it is the case, the heavy quark expansion seems to fail in the inclusive b decays except there are some special reasons to explain why the $1/m_Q^3$ terms become dorminant and the higher order terms $O(1/m_Q^4)$ are smaller. The nonspectator effects have recently been considered in ref. [12] and found to result in the following predictions

$$\frac{\tau(B^-)}{\tau(B^0)} = 1.11 \pm 0.02, \tag{1.2a}$$

$$\frac{\tau(B_s^0)}{\tau(B^0)} \approx 1, \tag{1.2b}$$

$$\frac{\tau(\Lambda_b)}{\tau(B^0)} \approx 0.99 - (\frac{f_B}{185\ \text{MeV}})^2(0.007 + 0.020\tilde{B})r \geq 0.98, \tag{1.2c}$$

where $\tilde{B}$ and r characterize the nonfactorization effects. Such a confliction has received wide attention. Authors in refs. [13,14] have also discussed the lifetime ratios and came to a similar result. Though the prediction for $\tau(B^-)/\tau(B^0)$ agrees with the current world average, those of $\tau(B_s^0)/\tau(B^0)$ and $\tau(\Lambda_b)/\tau(B^0)$ deviate somewhat from the central values of the world average. In the usual HQET, eq.(1.2b) is the final result because the only difference between B^0 and B^0_s decays lies in the different CKM matrix elements related in the nonspectator effects.

To understand the above problems, one of the attempts is to assume that the local duality may be violated in the nonleptonic inclusive decays. It was suggested in ref. [15] that a large $1/m_Q$ order correction in nonleptonic inclusive decays may exist and be simply described by replacing the heavy quark mass by the mass of the decaying hadron in the m^5 factor, i.e.,

$$\Gamma_{NL} \to \Gamma_{NL}(m_{H_b}/m_b)^5. \tag{1.3}$$

This assumption was further discussed in refs. [12,16,17]. This simple ansatz could not only resolve the lifetime ratio problem, but also provide the correct decay widths for the Λ_b baryon and the B mesons. But the charm counting may become much larger than the experimental data.

Secondly, it seems to have difficulties to simultaneously explain the semileptonic branching ratio B_{SL} and the charm counting n_c in B^0 decays. In general, when the charm counting is required to be near the experimental data, the predicted semileptonic branching ratio in the usual HQET is significantly larger than the experimental data. It has been shown in ref. [18] that for $m_b/2 \leq \mu \leq 2m_b$ and $m_b = 4.8 \pm 0.2$GeV

$$\begin{aligned} B_{SL} &= (12.0 \pm 0.7 \pm 0.5 \pm 0.2^{+0.9}_{-1.2})\%, &\text{(1.4a)}\\ n_c &= 1.24 \mp 0.05 \pm 0.01, &\text{(1.4b)}\\ \bar{B}_{SL} &= (11.3 \pm 0.6 \pm 0.7 \pm 0.2^{+0.9}_{-1.7})\%, &\text{(1.4c)}\\ \bar{n}_c &= 1.30 \mp 0.03 \pm 0.03 \pm 0.01, &\text{(1.4d)} \end{aligned}$$

where the first result is for the OS scheme and the second one for $\overline{MS}$ scheme. After considering the spectator effects, the results has been found to be improved [13]

$$B_{SL} = \begin{cases} 12.0 \pm 1.0\%; & \mu = m_b, \\ 10.9 \pm 1.0\%; & \mu = m_b/2, \end{cases} \tag{1.5a}$$

$$n_c = \begin{cases} 1.20 \mp 0.06; & \mu = m_b, \\ 1.21 \mp 0.06; & \mu = m_b/2. \end{cases} \tag{1.5b}$$

It is seen that the uncertainties in the two quantities are anti-correlated. One may compare them with the world average [20]

$$\begin{aligned} Br(b \to ce\bar{\nu}) &= 10.48 \pm 0.5\%, &\text{(1.6a)}\\ n_c &= 1.17 \pm 0.04. &\text{(1.6b)} \end{aligned}$$

Thirdly, there is an inconsistent picture between the Luke's theorem [19] for the exclusive heavy hadron decays and the Chay-Georgi-Grinstein theorem [3] for the inclusive heavy hadron decays. Luke's theorem tells us that the $1/m_Q$ order corrections are absent if one uses the meson mass to normalize the weak matrix elements in the exclusive heavy to heavy transitions at zero recoil. While according to the Chay-Georgi-Grinstein theorem, $1/m_Q$ order corrections are absent only when the quark mass is used. Consequently, the prediction of total decay width strongly depends on the value of quark mass.

These problems may arise from the simple treatment in the usual HQET, where the bound state effects and hadronization have not been taken into account. This may be a strong indication for a necessity of developing a new formulation of effective theory to incorporate these effects. For this purpose, we are going to devote in this paper the investigation of the inclusive bottom hadron decays and to provide better understanding on the problems mentioned above within the framework of the new formulation of HQEFT [2]. In section II, we briefly describe the framework of the new formulation of HQEFT. In section III, we apply it to the inclusive bottom hadron decays $H_b \to X_c \ell \nu$ and present a general formulation for $b \to c$ transitions via the heavy quark expansion. In section IV, we investigate the inclusive decays of B^0 and B_s^0 mesons as well as Λ_b baryon by providing detailed numerical results for their semileptonic branching ratios and lifetime ratios as well as the charm countings n_c. It is interesting to see that the results obtained by using the new formulation of HQEFT are consistent with the experimental data. In particular, the new formulation of HQEFT allows us to simply clarify the well known ambiguity of using the quark mass or hadron mass in the inclusive heavy hadron decays. As a consequence, the CKM matrix element $|V_{cb}|$ is well determined from the semileptonic decay rate. Conclusions and remarks are presented in the last section.

II. NEW FORMULATION OF HQEFT

For completeness, we present in this section a brief description on the framework of the new formulation of HQEFT [1,2]. Let us begin with the effective Lagrangian for a heavy quark

$$\begin{aligned}
\mathcal{L}_{eff} &= \bar{Q}_v^{(+)} iv\cdot D Q_v^{(+)} + \frac{1}{m_Q}\bar{Q}_v^{(+)}(i\not{D}_\perp)^2 Q_v^{(+)} \\
&\quad -\frac{1}{2m_Q^2}\bar{Q}_v^{(+)} i\not{D}_\perp iv\cdot D i\not{D}_\perp Q_v^{(+)} \qquad (2.1) \\
&\quad +\frac{1}{(2m_Q)^2}\bar{Q}_v^{(+)}(i\not{D}_\perp)^2 \frac{1}{iv\cdot D}(i\not{D}_\perp)^2 Q_v^{(+)} + O(1/m_Q^3) \qquad (2.2) \\
&= \mathcal{L}_{eff}^{(0)} + \mathcal{L}_{eff}^{(1/m_Q)},
\end{aligned}$$

where $\overleftarrow{D}_\mu$ and $\not{D}_\perp^\mu$ are defined as

$$\chi\overleftarrow{D}_\mu \equiv \partial_\mu\chi + i\chi g A_\mu^a T^a; \quad \not{D}_\perp^\mu \equiv (g^{\mu\nu} - v^\mu v^\nu) D_\nu$$

and $\mathcal{L}_{eff}^{(0)} = \bar{Q}_v^{(+)} iv\cdot D Q_v^{(+)}$ denotes the leading term and $\mathcal{L}_{eff}^{(1/m_Q)}$ represents the terms suppressed by the powers of $1/m_Q$. And the representation of left-handed current $J^\mu = \bar{Q}'\Gamma^\mu Q$ is

$$\begin{aligned}
J_{eff}^\mu &= e^{i(m_{Q'}v' - m_Q v)\cdot x}\bar{Q}_{v'}^{\prime(+)}\{\Gamma^\mu + \frac{1}{2m_{Q'}}(i\overleftarrow{\not{D}}_\perp{}')^2 \frac{1}{-iv'\cdot\overleftarrow{D}}\Gamma^\mu \\
&\quad +\frac{1}{2m_Q}\Gamma^\mu\frac{1}{iv\cdot D}(i\not{D}_\perp)^2\} Q_v^{(+)} + O(1/m_Q^2), \qquad (2.3)
\end{aligned}$$

with $\Gamma^\mu = \gamma^\mu \frac{1-\gamma^5}{2}$ and $D'^\mu_\perp = (g^{\mu\nu} - v'^\mu v'^\nu) D_\nu$.

In order to exhibit a manifest spin-flavor symmetry in the HQEFT, it is useful to introduce a hadron state $|H_v>$ corresponding to the effective heavy quark field $Q_v^{(+)}$. The hadron state $|H_v>$ is related to the state $|H>$ of heavy hadron H by the following equation

$$< H'|J^\mu|H >= \sqrt{\frac{m_{H'} m_H}{\bar{\Lambda}_{H'} \bar{\Lambda}_H}} < H'_{v'}|J^\mu{}_{eff} e^{i\int d^4x (\mathcal{L}_{eff}^{(1/m_Q)} + \mathcal{L}_{eff}^{(1/m_{Q'})})} |H_v >, \tag{2.4}$$

with m_H being the mass of the heavy hadron H and $\bar{\Lambda}_H = m_H - m_Q$. The factor m_H comes from the standard normalization

$$< H(p')|H(p) >= 2p^0 (2\pi)^3 \delta^3(\mathbf{p} - \mathbf{p}'). \tag{2.5}$$

Where the factor $\sqrt{\bar{\Lambda}_{H'} \bar{\Lambda}_H}$ is introduced to make the normalization concerning the hadron state $|H_v>$ to be independent of the heavy flavor, i.e.,

$$< H_v|\bar{Q}_v^{(+)} \gamma^\mu Q_v^{(+)}|H_v > \ = 2\bar{\Lambda} v^\mu, \tag{2.6}$$

with

$$\bar{\Lambda} = \lim_{m_Q \to \infty} \bar{\Lambda}_H.$$

Expanding the right-hand side of eq.(2.4), to the order of $1/m_Q$, we have

$$\begin{aligned} < H'|J^\mu|H > \ = & \sqrt{\frac{m_H m_{H'}}{\bar{\Lambda}_H \bar{\Lambda}_{H'}}} < H'_{v'}|\bar{Q}'^{(+)}_{v'} \{\Gamma^\mu - \frac{1}{2m_{Q'}} (i\overleftarrow{\not{D}'_\perp})^2 \frac{1}{-iv' \cdot \overleftarrow{D}} \Gamma^\mu \\ & - \frac{1}{2m_Q} \Gamma^\mu \frac{1}{iv \cdot D} (i\not{D}_\perp)^2 + O(\frac{1}{m^2_{Q^{(\prime)}}}) \} Q^{(+)} |H_v > . \end{aligned} \tag{2.7}$$

In general, a heavy quark in a hadron cannot truly be on-shell due to strong interactions among heavy quark and light quark as well as soft gluons. The off-shellness of heavy quark in the hadron is characterized by a residual momentum k. The total momentum P_Q of the heavy quark in a hadron may be written as: $P_Q = m_Q v + k$. In the usual HQET one mainly deals with the heavy quark and treats the light quark as a spectator, which does not affect the properties of heavy hadrons to a large extent (except the effects of weak annihilation and Pauli interference). However, since the light degrees of freedom affect the character of heavy hadrons, it may be this simple treatment that meets difficulties in explaining the lifetime differences between B^0 and B_s^0 as well as between Λ_b and B as mentioned in the introduction. To take the effects of light degrees into account and not to deal with hadronization directly, we will adopt an alternative picture, namely the residual momentum k of the heavy quark within a hadron is considered to comprise the contributions of the light degrees of freedom. With this picture the heavy quark may be regarded as a 'dressed heavy quark', thus the heavy hadron containing a single heavy quark may be more reliable to be considered as a dualized particle of a 'dressed heavy quark'. Thus the momentum P_H of a hadron H is decomposed into

$$P_H = m_Q v + k + k', \tag{2.8}$$

with k' being the momentum depending on the heavy flavor and suppressed by $1/m_Q$. With this picture, the momentum of the 'dressed heavy quark' inside the hadron is given by

$$P_Q = \lim_{m_Q \to \infty} P_H. \tag{2.9}$$

Hence

$$P_H^2 = m_H^2 = m_Q^2 + 2m_Q v \cdot (k + k') + k'^2 + k^2 + 2k \cdot k'. \tag{2.10}$$

Then

$$\bar{\Lambda}_H \equiv m_H - m_Q = \frac{v \cdot (k + k')}{1 + \bar{\Lambda}/2m_Q} + \frac{(k + k')^2}{2m_Q(1 + \bar{\Lambda}/2m_Q)}, \tag{2.11}$$

and

$$\bar{\Lambda} = \lim_{m_Q \to \infty} \bar{\Lambda}_H = \lim_{m_Q \to \infty} v \cdot (k + k') = v \cdot k. \tag{2.12}$$

In terms of the operator formulation, it implies that

$$< H_v|\bar{Q}_v^{(+)} iv \cdot D Q_v^{(+)}|H_v >= \bar{\Lambda} < H_v|\bar{Q}_v^{(+)} Q_v^{(+)}|H_v > . \tag{2.13}$$

Thus to simplify the evaluation of the matrix elements, one may approximately replace the propagator $1/iv \cdot D$ by $1/\bar{\Lambda}$

$$\begin{aligned} \kappa_1 &\equiv - < H_v|\bar{Q}_v^{(+)} D_\perp^2 Q_v^{(+)}|H_v > /(2\bar{\Lambda}), \\ \kappa_2 &\equiv < H_v|\bar{Q}_v^{(+)} g\sigma_{\mu\nu} G^{\mu\nu} Q_v^{(+)}|H_v > /(4 d_H \bar{\Lambda}). \end{aligned} \tag{2.14}$$

where $d_H = -3$ for pseudo scalar mesons, $d_H = 1$ for vector mesons and $d_H = 0$ for ground state heavy baryons.

With this approximation, the mass formulae can be simply given in terms of the effective Lagrangian $\mathcal{L}_{eff}^{(1/m_Q)}$

$$\begin{aligned} m_H &= m_Q + \bar{\Lambda} - \frac{< H_v|\bar{Q}_v^{(+)} (i\not{D}_\perp)^2 Q_v^{(+)}|H_v >}{2\bar{\Lambda} \cdot m_Q} + O(\frac{1}{m_Q^2}) \\ &\approx m_Q + \bar{\Lambda} - < H_v|\mathcal{L}_{eff}^{(1/m_Q)}|H_v > /(2\bar{\Lambda}), \end{aligned} \tag{2.15}$$

which can be reexpressed as follows by using eq.(2.14)

$$m_H \approx m_Q + \bar{\Lambda} - \frac{\kappa_1}{m_Q} + \frac{d_H \kappa_2}{m_Q} + O(\frac{1}{m_Q^2}). \tag{2.16}$$

It is useful to define the mass of 'dressed heavy quark' as

$$\hat{m}_Q \equiv \lim_{m_Q \to \infty} m_H = m_Q + \bar{\Lambda}, \tag{2.17}$$

which can be expressed in terms of the hadron mass

$$\begin{aligned} \hat{m}_Q &= m_H + \frac{\kappa_1 - d_H \kappa_2}{m_Q} + O(\frac{1}{m_Q^2}) \\ &= m_H + \frac{\kappa_1 - d_H \kappa_2}{m_H} + O(\frac{1}{m_H^2}). \end{aligned} \tag{2.18}$$

III. DYNAMICS OF INCLUSIVE DECAYS

The techniques for inclusive decays of heavy hadrons in the framework of effective theory was developed in the early years of this decade. Here we shall extend the method of ref. [6] to the framework of new formulation of HQEFT. Let us first briefly recall the basic formulae for the description of inclusive semileptonic decay

$$H(P_H = m_H v) \to X_c(P_X) + \ell(p) + \bar{\nu}_\ell(p'); \quad q = p + p',$$

which is mediated by the effective Hamiltonian

$$\mathcal{H}_{eff} = \frac{4G_F}{\sqrt{2}} V_{cb} \bar{b} \Gamma^\mu c \bar{\nu}_\ell \Gamma_\mu \ell = \frac{4G_F}{\sqrt{2}} V_{cb} J^\mu J_{\ell\mu}. \tag{3.1}$$

To calculate the decay rates, it is useful to introduce the hadronic tensor

$$W^{\mu\nu} = \frac{1}{2m_H} (2\pi)^3 \sum_{X_c} \delta^4(P_{H_b} - q - P_X) < H|J^{\mu\dagger}|X_c >< X_c|J^\nu|H >, \tag{3.2}$$

which can be expanded in terms of five form factors when one averages the spin of initial and final states

$$W^{\mu\nu} = -g^{\mu\nu} W_1 + v^\mu v^\nu W_2 - i\epsilon^{\mu\nu\alpha\beta} v_\alpha q_\beta W_3 + q^\mu q^\nu W_4 + (q^\mu v^\nu + q^\nu v^\mu) W_5. \tag{3.3}$$

For massless lepton pair, W_4 and W_5 will not contribute to the decay width

$$\frac{d\Gamma}{dq^2 dE_e dE_\nu} = \frac{G_F^2 V_{cb}^2}{2\pi^3} \Theta(E_e - q^2/4E_\nu) \{ q^2 W_1 + \frac{1}{2}(E_e E_\nu - q^2) W_2 + q^2 (E_e - E_\nu) W_3 \}. \tag{3.4}$$

To evaluate the form factors W_i, let us consider the time ordered product

$$\begin{aligned} T^{\mu\nu} &= -\frac{i}{2m_H} \int d^4x e^{-iq\cdot x} < H|\mathcal{T}\{J^{\mu\dagger}(x) J^\nu(0)\}|H > \\ &= -g^{\mu\nu} T_1 + v^\mu v^\nu T_2 - i\epsilon^{\mu\nu\alpha\beta} v^\alpha q^\beta T_3 + q^\mu q^\nu T_4 + (v^\mu q^\nu + v^\nu q^\mu) T_5. \end{aligned} \tag{3.5}$$

It is known that the form factors W_i are related to T_i via

$$W_i = -\frac{1}{\pi} Im T_i. \tag{3.6}$$

Explicitly, the quark matrix element in eq.(3.5) is given by

$$\frac{1}{2m_H} < H|\bar{b} \frac{\gamma^\mu (m_b \not{v} + \not{k} - \not{q}) \gamma^\nu}{(m_b v + k - q)^2 - m_c^2 + i\epsilon} \frac{1 - \gamma_5}{2} b|H > . \tag{3.7}$$

Here $m_b v + k - q = P_c$ is the momentum of charm quark.

Within the framework of new formulation of HQEFT, as we have discussed above, the residual momentum of the bottom quark in the hadron is given by $v \cdot k \sim \bar{\Lambda}$. For ensuring the leading term to have the largest contribution, we may expand eq.(3.7) in terms of the small subtracted momentum

$$k - v < v \cdot k > .$$

Here

$$< v\cdot k > \equiv < H|\bar{b}e^{-im_b \not{v}\cdot x} iv\cdot D e^{im_b \not{v}\cdot x} b|H > /2m_H = \bar{\Lambda}.$$

To evaluate the matrix element eq.(3.7), we need to know the following relevant matrix elements

$$\frac{1}{2m_H} < H|\bar{b}b|H > = 1, \tag{3.8a}$$

$$\frac{1}{2m_H} < H|\bar{b}e^{-im_b \not{v}\cdot x}\frac{1}{2}(iD_\perp^\kappa iD_\perp^\tau + iD_\perp^\tau iD_\perp^\kappa)e^{im_b \not{v}\cdot x} b|H > = A(g^{\kappa\tau} - v^\kappa v^\tau), \tag{3.8b}$$

$$\frac{1}{12m_H} < H|\bar{b}g\sigma_{\mu\nu}G^{\mu\nu}b|H > = N_b. \tag{3.8c}$$

Up to the leading order in $1/m_b$, we have, from eqs.(2.13) and (2.14), the following results for A and N_b

$$A = \frac{\kappa_1}{3}, \tag{3.9a}$$

$$N_b = \frac{d_H \kappa_2}{3}. \tag{3.9b}$$

The matrix element concerning one gluon in eq.(3.5) has the form

$$< H|\bar{b}\frac{g}{2}G^{\alpha\beta}\epsilon_{\alpha\beta\kappa\sigma}\frac{(m_b v + k - q)^\kappa}{2m_H}\frac{g^{\mu\sigma}\gamma^\nu + g^{\nu\sigma}\gamma^\mu - g^{\mu\nu}\gamma^\sigma + i\epsilon^{\mu\sigma\nu\tau}}{[(m_b v + k - q)^2 - m_c^2 + i\epsilon]^2}\Gamma^\tau b|H > . \tag{3.10}$$

Expanding eqs.(3.7) and (3.10), one can extract the form factors T_i. Using the relation eq.(3.6), one then obtains W_i. Here we shall only list the final results for W_i

$$\begin{aligned} W_1 = {} & \frac{1}{2}(\hat{m}_b - E_e - E_\nu)\delta(z) + \{\frac{1}{2}(3A + N_b)(\hat{m}_b - E_e - E_\nu)\}\delta'(z) \\ & + A(\hat{m}_b - E_e - E_\nu)\{q^2 - (E_e + E_\nu)^2\}\delta''(z), \end{aligned} \tag{3.11a}$$

$$\begin{aligned} W_2 = {} & \hat{m}_b\delta(z) + \{\hat{m}_b(3A - N_b) + 2A(E_e + E_\nu)\}\delta'(z) \\ & + 2\hat{m}_b A\{q^2 - (E_e + E_\nu)^2\}\delta''(z), \end{aligned} \tag{3.11b}$$

$$W_3 = \frac{1}{2}\delta(z) - \{\frac{1}{2}(5A + N_b)\}\delta'(z) + A\{q^2 - (E_e + E_\nu)^2\}\delta''(z), \tag{3.11c}$$

with $z = (\hat{m}_b v - q)^2 - m_c^2 = \hat{m}_b^2 + q^2 - 2\hat{m}_b E_e - 2\hat{m}_b E_\nu - m_c^2$.

Substituting eq.(3.11) into eq.(3.4) and integrating out the variables E_ν and q^2, one yields the inclusive lepton spectrum

$$\begin{aligned} \frac{1}{\hat{\Gamma}_0}\frac{d\Gamma}{dy} = {} & -2\rho^2(3-\rho) + \frac{4\rho^3}{(1-y)^3} - \frac{6\rho^2(1+\rho)}{(1-y)^2} + \frac{12\rho^2}{1-y} + 6(1-\rho)y^2 - 4y^3 \\ & + \frac{A}{\hat{m}_b^2}\{-6\rho^3 + 12\rho^2 - \frac{24\rho^3}{(1-y)^5} + \frac{6\rho^2(3+5\rho)}{(1-y)^4} - \frac{12\rho^2}{(1-y)^3} - \frac{18\rho^2}{(1-y)^2} + 6y^2\} \\ & + \frac{N_b}{\hat{m}_b^2}\{-6\rho(2+\rho) - \frac{12\rho^2}{(1-y)^3} + \frac{6\rho(2+3\rho)}{(1-y)^2} - 24\rho y - 18y^2\}, \end{aligned} \tag{3.12}$$

with

$$\hat{\Gamma}_0 \equiv \frac{G_F^2 \hat{m}_b^5 V_{cb}^2}{192\pi^3} = \frac{G_F^2 m_H^5 V_{cb}^2}{192\pi^3} \left(1 + \frac{\kappa_1 - d_H \kappa_2}{m_H^2} + O(\frac{1}{m_H^3})\right)^5$$
$$\equiv \Gamma_0 \left(1 + \frac{\kappa_1 - d_H \kappa_2}{m_H^2} + O(\frac{1}{m_H^3})\right)^5 . \qquad (3.13)$$

Where y and ρ are rescaled variables defined as

$$y \equiv 2E_e/\hat{m}_b, \quad \rho \equiv m_c^2/\hat{m}_b^2.$$

The kinetic region for y is

$$0 \leq y \leq \frac{P_b^2 - P_c^2}{P_b^2} - \frac{E_\nu}{2v \cdot P_b}. \qquad (3.14)$$

Treating charm quark c as heavy quark and using $v \cdot k = \bar{\Lambda}$, one has

$$0 \leq y \leq 1 - \hat{\rho}, \quad \hat{\rho} \equiv \hat{m}_c^2/\hat{m}_b^2. \qquad (3.15)$$

Thus the total decay width for $b \to ce\bar{\nu}$ is found to be

$$\frac{\Gamma}{\hat{\Gamma}_0} = -(1-\hat{\rho})^4 + 2(1-\hat{\rho})^3(1-\rho) - 2\rho^2(1-\hat{\rho})(3-\rho)$$
$$-2\rho^3(1 - \frac{1}{\hat{\rho}^2} + 6\rho^2(1+\rho)(1-\frac{1}{\hat{\rho}}) - 12\rho^2 \ln\hat{\rho}$$
$$+2\frac{A}{\hat{m}_b^2 \hat{\rho}^4}(1-\hat{\rho})^3\{\hat{\rho}^4 + \hat{\rho}\rho^2(3-4\rho) + 3\hat{\rho}^2\rho^2(2-\rho) - 3\rho^3\}$$
$$-\frac{6N_b}{\hat{m}_b^2 \hat{\rho}^2}(1-\hat{\rho})^3(\hat{\rho}-\rho)^2. \qquad (3.16)$$

Similarly, we find that $b \to c$ transitions have the following general forms

$$\frac{1}{\hat{\Gamma}_0}\Gamma(b \to c\ell\nu) = \eta_{c\ell}(\rho, \rho_\ell, \mu)\{I_0(\rho, \rho_\ell, \hat{\rho}) + I_1(\rho, \rho_\ell, \hat{\rho})\frac{A}{\hat{m}_b^2} + I_2(\rho, \rho_\tau, \hat{\rho})\frac{N_b}{\hat{m}_b^2}\}, \qquad (3.17a)$$

$$\frac{1}{\hat{\Gamma}_0}\Gamma(b \to cud') = 3\eta_{cu}(\rho, \mu)\{I_0(\rho, 0, \hat{\rho}) + I_1(\rho, 0, \hat{\rho})\frac{A}{\hat{m}_b^2}$$
$$+I_2(\rho, 0, \hat{\rho})\frac{N_b}{\hat{m}_b^2}\} - (c_+^2(\mu) - c_-^2(\mu))6(1-\rho)^3\kappa_2/\hat{m}_b^2, \qquad (3.17b)$$

$$\frac{1}{\hat{\Gamma}_0}\Gamma(b \to ccs') = 3\eta_{cc}(\rho, \mu)\{I_0(\rho, \rho, \hat{\rho}) + I_1(\rho, \rho, \hat{\rho})\frac{A}{\hat{m}_b^2}$$
$$+I_2(\rho, \rho, \hat{\rho})\frac{N_b}{\hat{m}_b^2}\} - (c_+^2(\mu) - c_-^2(\mu))I_3(\rho, \rho, 0)6\kappa_2/\hat{m}_b^2, \qquad (3.17c)$$

with $s' = dV_{cd} + sV_{cs}$ and $d' = dV_{ud} + sV_{us}$ and $\rho_\ell \equiv m_\ell^2/\hat{m}_b^2$. The η functions arise from QCD radiative corrections. The one-loop results have been calculated in refs. [18,21,22]. Here we will use the two-loop results in refs. [23,24] and adopt the reference value $m_c/m_b = 0.3$, which leads to the following results

$$\eta_{ce} = 1 - 0.53\alpha_s(\mu) - 1.53\alpha_s(\mu)^2, \tag{3.18a}$$

$$\eta_{c\tau} = 1 - 0.44\alpha_s(\mu) - 1.44\alpha_s(\mu)^2, \tag{3.18b}$$

$$\eta_{cu} = 1 + 0.95\alpha_s(\mu) + \frac{\alpha_s(\mu)^2}{\pi^2}[33.03 + 3.34\ln(m_W/m_b) + 4\ln(m_W/m_b)^2], \tag{3.18c}$$

$$\eta_{cc} = 1 - 0.21\alpha_s(\mu) + \frac{\alpha_s(\mu)^2}{\pi^2}[-9.99 + 7.17\ln(m_W/m_b) + 4\ln(m_W/m_b)^2]. \tag{3.18d}$$

The functions $I_i(x,y,0)$ are phase space factors at tree level. Their explicit forms are

$$\begin{aligned} I_0(x,y,z) = {} & \sqrt{\kappa}\{1 - z - z^2 + z^3 - 2\,z^2\,x + 2\,x^3 - 2\,x\,(-1+y)^2 \\ & - \frac{6\,x^2\,(-1+y)^2}{z} + \frac{2\,x^3\,(-1+y)^2}{z^2} - 7\,y - z^2\,y - 7\,y^2 - z\,y^2 \\ & + y^3 + 4\,z\,x\,(1+y) - 6\,x^2\,(1+y) - \frac{4\,x^3\,(1+y)}{z}\} \\ & + 12x^2\ln\frac{(1+z-y+\sqrt{\kappa})^2}{4z} + 12y^2\ln\frac{(1-z+y+\sqrt{\kappa})^2}{4y} \\ & - 12x^2y^2\ln\frac{(1-z-y+\sqrt{\kappa})^2}{4zy}, \end{aligned} \tag{3.19a}$$

$$\begin{aligned} I_1(x,y,z) = {} & \frac{2\sqrt{\kappa}}{z^4}\{z^6 - 3\,x^3\,(-1+y)^4 - 2\,z^5\,(1+y) + z^3\,x^2\,(-9 + 2\,x \\ & - 18\,y + 12\,x\,y + 39\,y^2) + z^4\,[-3\,x^3 + (-1+y)^2 + 3\,x^2\,(2+3\,y)] \\ & + z\,x^2\,(-1+y)^2\,[3\,(-1+y)^2 + 2\,x\,(1+6\,y)] \\ & + z^2\,x^2\,[-15\,(-1+y)^2\,y + x\,(2 + 8\,y - 18\,y^2)]\} \\ & + 72x^2y^2\ln\frac{(1-z-y+\sqrt{\kappa})^2}{4zy}, \end{aligned} \tag{3.19b}$$

$$I_2(x,y,z) = -6(x-z)^2[z^2 + (1-y)^2 - 2z(1+y)]^{\frac{3}{2}}, \tag{3.19c}$$

$$I_3(x,x,0) = \sqrt{1-4x}(1 + \frac{x}{2} + 3x^2) - 3x(1-2x^2)\ln\frac{1+\sqrt{1-4x}}{1-\sqrt{1-4x}}, \tag{3.19d}$$

with $\kappa = (1+y-z)^2 - 4y$. Note that for $z = x$, I_0 agrees with that in refs. [6,9].

Before ending this section, we would like to address the following points: Firstly, the mass entering into the factor $\hat{\Gamma}_0$ is neither the heavy quark mass m_b nor the hadron mass m_H, it is the so-called 'dressed heavy quark' mass $\hat{m}_b = m_b + \bar{\Lambda} = m_{H_b}(1 + O(1/m_{H_b}^2))$, which differs from the hadron mass by terms suppressed by $1/m_b$. Secondly, paralleling to Luke's theorem, there is no $1/m_b$ order correction when the hadron mass is used. Thus the uncertain parameters, i.e., the bottom quark mass m_b and binding energy $\bar{\Lambda}$, do not enter separately into the expression of decay widths. Furthermore, one may notice that for the final charm quark, both charm quark mass m_c and 'dressed charm quark' mass $\hat{m}_c$ have entered into the phace space factors of the general formulation of the decay rates. Where the quark mass m_c comes from the propagator of charm quark, and the 'dressed charm quark' mass $\hat{m}_c$ arises from the momentum of charm quark inside the hadron, i.e., $P_c = P_b - q = \hat{m}_b v - q = \hat{m}_c v'$, for a parallel treatment to the bottom quark inside the hadron. These features enable us to determine the lifetime ratios, semileptonic branching ratios and even the lifetimes of bottom hadrons in rather accuracy.

IV. NUMERICAL RESULTS AND DISCUSSIONS

In our numerical calculations, we shall neglect the contributions from penguin diagrams and other rare decays and incorporate the two-loop radiative corrections in refs. [23,24] to calculate the values of observable parameters in b decays. Such an approximation causes errors no more than 2%. Thus the total decay width for bottom hadrons are given by

$$\Gamma_H^t = \frac{1}{\tau(H)} \simeq \Gamma(b \to cud') + \Gamma(b \to ccs') + \sum_\ell \Gamma(b \to c\ell\bar{\nu}) + \Gamma(b \to u). \tag{4.1}$$

The semileptonic branching ratios are defined as

$$B_{SL}(H) \equiv Br(H_b \to X_c e\bar{\nu}) = \frac{\Gamma(H_b \to X_c e\bar{\nu})}{\Gamma_H^t},$$

$$B_\tau(H) \equiv Br(H_b \to X_c \tau\bar{\nu}) = \frac{\Gamma(H_b \to X_c \tau\bar{\nu})}{\Gamma_H^t}.$$

Other two ratios concerning the charm counting

$$n_c(H) = 1 + \frac{\Gamma(H_b \to X_{c\bar{c}})}{\Gamma_H^t} - \frac{\Gamma \underline{\text{ nocharm}}}{\Gamma_H^t}$$

and relative leptonic contributions between the τ decay $B_\tau(H)$ and the β decay $B_{SL}(H)$

$$R(H) \equiv \frac{B_\tau(H)}{B_{SL}(H)}.$$

Note that the formula given above are general and can be applied to bottom hadrons by just taking different binding energies $\bar{\Lambda}(H)$ for different bottom hadrons. We now discuss these observable quantities in detail below.

A. Input Parameters

The basic parameters involved are m_c, μ, κ_1 and κ_2. Using eq.(2.16) and the measured masses of the ground state heavy mesons and heavy baryons, one can find that κ_1 in the heavy baryon system is almost the same as the one in the heavy meson system. Thus besides the known masses [20]

$$\begin{aligned} m_{B^0} = 5.2792 \text{ GeV}; \quad m_{B^{*0}} = 5.3249 \text{ GeV}; \\ m_{D^+} = 1.8693 \text{ GeV}; \quad m_{D^{*+}} = 2.0100 \text{ GeV}; \\ m_{\Lambda_c} = 2.2849 \text{ GeV}; \quad m_{\Lambda_b} = 5.624 \text{ GeV}; \quad m_\tau = 1.777 \text{ GeV}, \end{aligned} \tag{4.2}$$

there are only four parameters m_c, μ, κ_1 and κ_2 in our calculations. Here κ_1 could be computed by QCD sum rules [26] or other phenomenological model [27]. It may also be extracted from fitting the meson spectra. Nevertheless, all the results remain suffering from large uncertainties. Here we shall use the most conservative range

$$0.3 \text{ GeV}^2 \leq -\kappa_1 \leq 0.7 \text{ GeV}^2. \tag{4.3}$$

For κ_2, the value extracted from the known $B - B^*$ mass splitting is quite stable

$$\kappa_2 = \frac{1}{8}(m^2_{B^{*0}} - m^2_{B^0}) = 0.06 \text{ GeV}^2, \tag{4.4}$$

which is accurate up to power correction of $\bar{\Lambda}/2m_b \sim 5\%$. The mass parameter m_c arising from the propagator in eqs.(3.7) and (3.10) shall be taken as an effective pole mass. Its value has a large range [20]

$$1.2 \text{ GeV} \leq m_c^{pole} \leq 1.9 \text{ GeV}.$$

Note that m_c^{pole} is different from the 'heavy quark mass' in the Lagrangian eq.(2.1) and the 'dressed heavy quark' mass in eq.(2.18). In the numerical calculations, we will restrict its range to be

$$1.55 \text{ GeV} \leq m_c^{pole} \leq 1.75 \text{ GeV}.$$

B. Lifetime Ratios

It is seen that, in the framework of new formulation of HQEFT, only 'dressed bottom quark' mass $\hat{m}_b$ enters into the decay rate, but both charm quark mass m_c and 'dressed charm quark' mass $\hat{m}_c$ appear in the phase space factors at tree level. So the lifetime ratio is not merely given by the ratio $[m_b + \bar{\Lambda}(H)]^5/[m_b + \bar{\Lambda}(H')]^5$. The μ, m_c and κ_1 dependences of the lifetime ratios $\tau(B_s^0)/\tau(B^0)$ and $\tau(\Lambda_b)/\tau(B^0)$ are plotted in Fig. 1. It is seen that the ratios are not sensitive to the energy scale μ. For a large range of parameters m_c and κ_1, the ratios only slightly change, but for large κ_1, the ratios become very sensitive to a large charm quark mass. The ratios as functions of m_c and κ_1 also exhibit some minimal points around which they change slowly. When taking 1.55 GeV $\leq m_c \leq$ 1.75 GeV and -0.7 GeV$^2 \leq \kappa_1 \leq -0.3$ GeV^2, we have

$$\frac{\tau(B_s^0)}{\tau(B^0)} = 0.94 \pm 0.04, \tag{4.5a}$$

$$\frac{\tau(\Lambda_b)}{\tau(B^0)} = 0.76 \pm 0.06, \tag{4.5b}$$

which show a good agreement with the experimental data [11].

C. Numerical Results in B^0 Decays

There is a puzzle in b decays in the usual HQET that the predicted semileptonic branching ratio is significantly greater than the experimental data. A large QCD enhancement of

$b \to c\bar{c}s$ was expected to suppress the value of the ratio, but it will lead to a much larger charm counting than the world average.

The theoretical predictions for B_{SL} and B_τ strongly depend on the energy scale μ, as shown in Fig. 2 and Fig. 3. At $\mu = m_b$ and $\mu = m_b/2$, we have

$$B_{SL} = \begin{cases} 11.98 \pm 0.60\%; & \mu = m_b, \\ 10.00 \pm 0.53\%; & \mu = m_b/2, \end{cases} \tag{4.6a}$$

$$B_\tau = \begin{cases} 3.23 \mp 0.21\%; & \mu = m_b, \\ 2.69 \mp 0.19\%; & \mu = m_b/2. \end{cases} \tag{4.6b}$$

Their uncertainties caused by the charm quark mass and κ_1 are anti-correlated. One may compare it with the usual HQET prediction as shown in Fig. 6a and 6b.

But their ratio R is not sensitive to the choice of the scale μ as shown in Fig. 4. As the function of m_c and κ_1, it shows a minimal point around which its variations become slow. For 1.55 GeV $\leq m_c \leq$ 1.75 GeV and -0.7 GeV$^2 \leq \kappa_1 \leq -0.3$ GeV^2, we have

$$0.24 \leq R \leq 0.32, \tag{4.7}$$

here the main uncertainties arise from the uncertainties of κ_1 and m_c. One may also compare it with the usual HQET prediction as shown in Fig. 6c. For $m_b = 4.7 \pm 0.1$ GeV, $a = 0.29 \pm 0.03$ and $\lambda_1 = -0.36$ GeV^2, one has $0.18 < R < 0.26$. This shows one of the differences between the framework of new formulation of HQEFT and the usual HQET. The CLEO data for B_{SL} and B_τ is [25]

$$B_{SL} = 10.5 \pm 0.5\%, \tag{4.8a}$$

$$B_\tau = 2.6 \pm 0.1\%, \tag{4.8b}$$

which leads to $R = 0.25 \pm 0.02$.

In analogous to R, n_c is not sensitive to μ, but it strongly depends on m_c and κ_1. The uncertainty in the theoretical prediction mainly arises from that of the value m_c as shown in Fig. 5. For 1.45 GeV $\leq m_c \leq$ 1.80 GeV and -0.7 GeV$^2 \leq \kappa_1 \leq -0.3$ GeV^2, the numerical result for n_c is

$$n_c = 1.18 \pm 0.06 \pm 0.03. \tag{4.9}$$

The uncertainties arising from the scale μ and Λ_{QCD} are not more than 0.03. Note that its m_c dependence shows a minimal point around which its variation is relatively slow. One may compare it with the usual HQET prediction as shown in Fig. 6c. For $\mu = m_b/2 - 2m_b$, $m_b = 4.7 \pm 0.1$ GeV, $a = 0.29 \pm 0.03$ and $\lambda_1 = -0.36$ GeV^2, one has $n_c = 1.22 - 1.34$.

From Figs. 2-5, one can see that there exists a common region for parameters μ, m_c and κ_1, so that all the quantities B_{SL}, B_τ, R and n_c are consistent with experimental data. When taking the following interesting region for the parameters

$$\begin{aligned} 1.55 \text{ GeV} \leq m_c \leq 1.75 \text{ GeV}, \\ 2.4 \text{ GeV} \leq \mu \leq 4.4 \text{ GeV}, \\ -0.7 \text{ GeV}^2 \leq \kappa_1 \leq -0.3 \text{ GeV}^2, \end{aligned} \tag{4.10}$$

and normalizing $B_{SL} = 10.48 \pm 0.50\%$ and $R = 0.26 \pm 0.02$, we have

$$n_c = 1.22 \mp 0.05. \tag{4.11}$$

which is consistent with the world average value. Note that a larger m_c value leads to a lower R and n_c but to a larger B_{SL} and B_τ.

In the usual HQET, up to $1/m_Q^2$ order and neglecting the 'spectator effect', however, there does not exist a common region for any choice of parameters, which makes all the quantities to be consistent with the experimental data. This can been explicitly seen from Fig. 6., where we have plotted four quantities B_{SL}, B_τ, R and n_c as the functions of the running energy scale μ and the mass ratio $a \equiv m_c/m_b$ with $m_b = 4.8 \pm 0.2$ GeV (Note that in the usual HQET, the b quark mass m_b, instead of the 'dressed heavy quark' mass $\hat{m}_b$, is the basic parameter). It is clear from Fig. 6 that there are two obstacles in the usual HQET to obtain a consistent fit. Firstly, a lower n_c value requires a larger mass ratio a and scale μ, while a higher scale will lead semileptonic branching ratio to be larger than the experimental bound, and a higher mass ratio will result in a lower value R.

D. Predictions for B_s^0 and Λ_b Decays

We now discuss more about B_s^0 and Λ_b decay. In the usual HQET, all the above observable quantities in the B^0, B_s^0 and Λ_b decays get the same results up to $1/m_b^2$ order. This is because the semileptonic decay rates of all the bottom hadrons are the same, thus the usual HQET leads to the following predictions up to $1/m_b^2$ order

$$\frac{B_{SL}(B_s^0)}{B_{SL}(B^0)} = \frac{\Gamma^t_{B^0}}{\Gamma^t_{B_s^0}} = \left(\frac{\tau(B_s^0)}{\tau(B^0)}\right)_{th} \approx 1, \tag{4.12a}$$

$$\frac{B_{SL}(B^-)}{B_{SL}(B^0)} = \frac{\Gamma^t_{B^0}}{\Gamma^t_{B^-}} = \left(\frac{\tau(B^-)}{\tau(B^0)}\right)_{th} \approx 1, \tag{4.12b}$$

$$\frac{B_{SL}(\Lambda_b)}{B_{SL}(B^0)} = \frac{\Gamma^t_{B^0}}{\Gamma^t_{\Lambda_b}} = \left(\frac{\tau(\Lambda_b)}{\tau(B^0)}\right)_{th} \approx 1. \tag{4.12c}$$

In the usual HQET, the lifetime ratio problems are expected to be solved through the nonleptonic decays. If it is the case, one should have the following consequences for the ratios of the semileptonic branching ratios

$$\frac{B_{SL}(B_s^0)}{B_{SL}(B^0)} = \left(\frac{\tau(B_s^0)}{\tau(B^0)}\right)_{exp} = 0.94 \pm 0.04, \tag{4.13a}$$

$$\frac{B_{SL}(B^-)}{B_{SL}(B^0)} = \left(\frac{\tau(B^-)}{\tau(B^0)}\right)_{exp} = 1.07 \pm 0.03, \tag{4.13b}$$

$$\frac{B_{SL}(\Lambda_b)}{B_{SL}(B^0)} = \left(\frac{\tau(\Lambda_b)}{\tau(B^0)}\right)_{exp} = 0.79 \pm 0.05. \tag{4.13c}$$

For the ratio R one has

$$R(B^0) \approx R(B^-) \approx R(B_s^0) \approx R(\Lambda_b). \tag{4.14}$$

One may compare it with the current experimental data [20]

$$
\begin{aligned}
B_{SL}(B^0) &= 10.48 \pm 0.5\%, && (4.15a)\\
B_{SL}(B^-) &= 10.18 \pm 0.5\%, && (4.15b)\\
B(B^0_s \to D^- \ell^+ \bar{\nu} \text{ anything}) &= 8.1 \pm 2.5\%, && (4.15c)\\
B(\Lambda_b \to \Lambda_c \ell \bar{\nu} \text{ anything}) &= 9.0^{+3.0}_{-2.8}\%. && (4.15d)
\end{aligned}
$$

In the framework of new formulation of HQEFT, as we have discussed in the previous sections, the picture for heavy hadron containing a single heavy quark is such that the heavy quark in the hadron is off-shell, its off-shellness is characterized by the binding energy $\bar{\Lambda}$. Consequently, the so-called 'dressed heavy quark' masses $\hat{m}_b$ and $\hat{m}_c$ enter into the phase space factors. For B^0_s and Λ_b decays, the differences of the phase space factors already appear at tree level. It is not difficult to obtain the following relations

$$
\begin{aligned}
&B_{SL}(B^0) < B_{SL}(B^0_s) < B_{SL}(\Lambda_b), && (4.16a)\\
&R(B^0) \lesssim R(B^0_s) < R(\Lambda_b), && (4.16b)\\
&n_c(B^0) > n_c(B^0_s) > n_c(\Lambda_b). && (4.16c)
\end{aligned}
$$

Compared eqs.(4.13) and eq.(4.14) with eq.(4.16), one may find the significant difference between the usual HQET and the new formulation. The more and more accurate experiments in the future will provide results of $\mathcal{B}_{SL}$ and R to test the usual HQET and the new formulation of HQEFT.

When normalizing $B_{SL}(B^0)$ to 9.98% ∼ 10.98% and $R(B^0)$ to 0.24 ∼ 0.28, we have

$$
\begin{aligned}
B_{SL}(B^0_s) &= 10.75 \pm 0.67\%; \qquad & B_{SL}(\Lambda_b) &= 11.09 \pm 0.91\%, && (4.17a)\\
\frac{B_{SL}(B^0_s)}{B_{SL}(B^0)} &= 1.02 \pm 0.02; \qquad & \frac{B_{SL}(\Lambda_b)}{B_{SL}(B^0)} &= 1.06 \pm 0.05, && (4.17b)\\
R(B^0_s) &= 0.28 \mp 0.02; \qquad & R(\Lambda_b) &= 0.32 \mp 0.02, && (4.17c)\\
n_c(B^0_s) &= 1.20 \pm 0.04; \qquad & n_c(\Lambda_b) &= 1.18 \pm 0.05. && (4.17d)
\end{aligned}
$$

E. $|V_{cb}|$ from Inclusive Decays

One of the most important applications is the determination of the CKM matrix element $|V_{cb}|$. From eq.(3.17a) the semileptonic decay rate is given by

$$
\Gamma(b \to c e \bar{\nu}) = \frac{G_F^2 \hat{m}_b^5 V_{cb}^2}{192\pi^3} \eta_{ce}(\rho, 0, \mu) \{ I_0(\rho, 0, \hat{\rho}) + I_1(\rho, 0, \hat{\rho}) \frac{A}{\hat{m}_b^2} + I_2(\rho, 0, \hat{\rho}) \frac{N_b}{\hat{m}_b^2} + \cdots \}, \quad (4.18)
$$

where the ellipse denotes higher order perturbative and non-perturbative corrections. It is explicitly seen that only the combination of $\bar{\Lambda}$ and m_b, i.e., $\hat{m}_b$, appears in the above equation and there is no $1/m_b$ corrections. At the same time, the $1/m_b$ power corrections to the decay width $\Gamma(b \to c e \bar{\nu})$ is as small as $-0.7 \sim 5\%$. Thus all the parameters in eq.(4.18) have clear physical meaning and the higher power corrections can be ignored. This allows us to extract $|V_{cb}|$ from fitting the experimental data. The value of $|V_{cb}|$ as function of m_c and μ is shown in Fig. 7. It is easy to find that the value of $|V_{cb}|$ has only a weak dependence on the energy scale μ and the charm quark mass m_c and κ_1 cause the main uncertainties for the prediction.

There exist maximal points around which its variation as the function of m_c and κ_1 becomes slow. The values of $|V_{cb}|$ around those points should be more reliable. It is interesting to note that once one normalizes the semileptonic branching ratio to the experimental data, $|V_{cb}|$ exhibits an interesting correlation with the ratio R. Its relation is shown in Fig. 7d. It is seen that the value of $|V_{cb}|$ increases linearly as R decreases. Including two-loop QCD corrections and fitting $Br(b \to ce\bar{\nu})$ to the measured data, we obtain

$$|V_{cb}| = 0.0388 \pm 0.0005_{exp} \pm 0.0012_{th}, \tag{4.19}$$

where the theoretical uncertainties arise from those of $R = 0.25 \pm 0.03$ and $\kappa_1 = -0.5 \pm 0.2$ GeV^2. The experimental uncertainty comes from the errors of B^0 lifetime. One may compare the above prediction for $|V_{cb}|$ in the new formulation of HQEFT with the one in the usual HQET which is plotted in Fig. 8.

V. CONCLUSIONS

We have presented, within the framework of new formulation of HQEFT, a general formulation for the inclusive decays of bottom hadrons via the heavy quark expansion. Such a general formulation has exhibited interesting features: The $1/m_b$ corrections are absent and the mass preliminarily entered into the general formulation is neither the bottom quark mass m_b nor heavy hadron mass m_H, it is the so-called 'dressed bottom quark' mass $\hat{m}_b$ that is actually the heavy hadron mass in the infinite quark mass limit, i.e., $\hat{m}_b = \lim_{m_b \to \infty} m_{H_b} = m_b + \bar{\Lambda} = m_{H_b}[1 + O(1/m_{H_b}^2)]$. Consequently, the decay rates of the bottom hadrons can be reexpressed in terms of the bottom hadron mass with the $1/m_b$ corrections remaining absent. This feature is now consistent with the expectation from Luke's theorem, since according to that theorem the sum of all the channels of exclusive b decays is free from the $1/m_b$ order corrections at zero recoil. Such a feature distinguishes from the one obtained by Chay-Georgi-Grinstein theorem within the framework of the usual HQET. The reason is that in the new formulation of HQEFT, Luke's theorem comes out automatically without the need of imposing the equation of motion $iv \cdot DQ_v^{(+)} = 0$. As a consequence, not only the lifetime ratio $\tau(\Lambda_b)/\tau(B^0)$ but also the ratio $\tau(B_s^0)/\tau(B^0)$ is in good agreement with the experimental data. At the same time, the results for the inclusive semileptonic branching ratio B_{SL}, the ratio R and the charm counting n_c are also consistent with the experimental data. In particular, the CKM matrix element $|V_{cb}|$ has been nicely extracted from the inclusive semileptonic decay rate, and the result well agrees with the one from the exclusive decays [1]. For $m_c = 1.75$ GeV and $\kappa_1 = -0.4$ GeV^2, we have

$$\frac{\tau(B_s^0)}{\tau(B^0)} = 0.95; \qquad \frac{\tau(\Lambda_b)}{\tau(B^0)} = 0.77, \tag{5.1a}$$

$$B_{SL}(B_s^0) = 10.82\%; \qquad B_{SL}(\Lambda_b) = 11.32\%, \tag{5.1b}$$

$$R(B^0) = 0.25; \quad R(B_s^0) = 0.27; \quad R(\Lambda_b) = 0.32, \tag{5.1c}$$

$$n_c(B^0) = 1.19; \quad n_c(B_s^0) = 1.17; \quad n_c(\Lambda_b) = 1.13, \tag{5.1d}$$

$$|V_{cb}| = 0.0398. \tag{5.1e}$$

If the nonspectator effects are taken into account, the charm counting will decrease further [13]. It is expected that more precise data for the B^0 decays and further test for the predictions on the B_s^0 and Λ_b systems will provide a useful check for the framework of new formulation of HQEFT at the level of higher order corrections.

ACKNOWLEDGMENTS

We would like to thank professor Y.B. Dai for useful discussions. This work was supported in part by the NSF of China under the grant No. 19625514.

Table. 1. The observable parameters in B^0 and B_s^0 as well as Λ_b decays with $\mu = m_b$.

m_c(GeV)	1.55			1.65			1.75			1.8		
κ_1(GeV^2)	-0.5	-0.6	-0.7	-0.4	-0.5	-0.6	-0.3	-0.4	-0.5	-0.2	-0.3	-0.4
$\frac{\tau(B_s^0)}{\tau(B^0)}$	0.903	0.915	0.946	0.913	0.931	0.969	0.923	0.945	0.987	0.922	0.938	0.970
$\frac{\tau(\Lambda_b)}{\tau(B^0)}$	0.701	0.703	0.724	0.718	0.729	0.764	0.738	0.757	0.804	0.743	0.757	0.793
$B_{SL}(B^0)(\%)$	11.62	11.56	11.46	12.04	11.92	11.74	12.41	12.20	11.94	12.68	12.44	12.16
$B_\tau(B^0)$	0.032	0.031	0.030	0.032	0.031	0.030	0.031	0.031	0.030	0.032	0.031	0.030
$\mathcal{R}(B^0)$	0.27	0.26	0.26	0.26	0.26	0.26	0.25	0.25	0.25	0.25	0.25	0.25
$n_c(B^0)$	1.21	1.22	1.23	1.18	1.19	1.21	1.16	1.17	1.19	1.14	1.16	1.18
$B_{SL}(B_s^0)(\%)$	11.68	11.66	11.63	12.19	12.12	12.04	12.68	12.55	12.39	12.99	12.82	12.63
$B_\tau(B_s^0)$	0.034	0.033	0.032	0.034	0.033	0.032	0.034	0.033	0.032	0.035	0.034	0.032
$\mathcal{R}(B_s^0)$	0.29	0.28	0.27	0.28	0.27	0.27	0.27	0.26	0.26	0.27	0.26	0.26
$n_c(B_s^0)$	1.21	1.21	1.21	1.17	1.18	1.18	1.14	1.15	1.16	1.12	1.13	1.14
$\frac{B_{SL}(B_s^0)}{B_{SL}(B^0)}$	1.00	1.01	1.02	1.01	1.02	1.03	1.02	1.03	1.04	1.02	1.03	1.04
$B_{SL}(\Lambda_b)(\%)$	11.73	11.72	11.73	12.37	12.35	12.33	13.08	13.02	12.96	13.49	13.40	13.32
$B_\tau(\Lambda_b)$	0.039	0.038	0.037	0.040	0.039	0.038	0.041	0.040	0.039	0.042	0.041	0.040
$\mathcal{R}(\Lambda_b)$	0.33	0.32	0.32	0.32	0.32	0.31	0.31	0.31	0.30	0.31	0.30	0.30
$n_c(\Lambda_b)$	1.21	1.21	1.21	1.16	1.16	1.17	1.11	1.12	1.12	1.08	1.09	1.10
$\frac{B_{SL}(\Lambda_b)}{B_{SL}(B^0)}$	1.01	1.01	1.02	1.03	1.04	1.05	1.05	1.07	1.09	1.06	1.08	1.10
$V_{cb}(10^{-2})$	3.63	3.73	3.72	3.68	3.73	3.74	3.75	3.79	3.77	3.76	3.81	3.82

Table. 2. The observable parameters in B^0 and B_s^0 as well as Λ_b decays with $\mu = m_b/2$.

m_c(GeV)	1.55			1.65			1.75			1.8		
κ_1(GeV^2)	-0.5	-0.6	-0.7	-0.4	-0.5	-0.6	-0.3	-0.4	-0.5	-0.2	-0.3	-0.4
$\frac{\tau(B_s^0)}{\tau(B^0)}$	0.904	0.917	0.949	0.916	0.934	0.973	0.927	0.949	0.993	0.927	0.943	0.976
$\frac{\tau(\Lambda_b)}{\tau(B^0)}$	0.706	0.708	0.731	0.725	0.737	0.774	0.748	0.769	0.818	0.755	0.770	0.809
$B_{SL}(B^0)$(%)	9.53	9.47	9.38	9.93	9.81	9.64	10.28	10.08	9.83	10.53	10.30	10.03
$B_\tau(B^0)$	0.026	0.026	0.025	0.027	0.026	0.025	0.026	0.026	0.025	0.027	0.026	0.025
$\mathcal{R}(B^0)$	0.28	0.27	0.27	0.27	0.26	0.26	0.26	0.25	0.26	0.26	0.25	0.25
$n_c(B^0)$	1.24	1.25	1.26	1.21	1.22	1.23	1.18	1.20	1.22	1.16	1.18	1.20
$B_{SL}(B_s^0)$(%)	9.60	9.58	9.55	10.08	10.01	9.93	10.55	10.41	10.26	10.84	10.68	10.49
$B_\tau(B_s^0)$	0.028	0.027	0.027	0.029	0.028	0.027	0.029	0.028	0.027	0.030	0.028	0.027
$\mathcal{R}(B_s^0)$	0.29	0.29	0.28	0.29	0.28	0.27	0.27	0.27	0.26	0.27	0.27	0.26
$n_c(B_s^0)$	1.24	1.24	1.24	1.20	1.20	1.21	1.16	1.17	1.19	1.14	1.15	1.17
$\frac{B_{SL}(B_s^0)}{B_{SL}(B^0)}$	1.01	1.01	1.02	1.02	1.02	1.03	1.03	1.03	1.04	1.03	1.04	1.05
$B_{SL}(\Lambda_b)$(%)	9.69	9.68	9.68	10.30	10.27	10.25	10.97	10.91	10.85	11.38	11.29	11.20
$B_\tau(\Lambda_b)$	0.032	0.032	0.031	0.034	0.033	0.032	0.035	0.034	0.033	0.036	0.035	0.034
$\mathcal{R}(\Lambda_b)$	0.34	0.33	0.32	0.33	0.32	0.32	0.32	0.31	0.30	0.31	0.31	0.30
$n_c(\Lambda_b)$	1.23	1.24	1.24	1.19	1.19	1.19	1.13	1.14	1.14	1.10	1.11	1.12
$\frac{B_{SL}(\Lambda_b)}{B_{SL}(B^0)}$	1.02	1.02	1.03	1.04	1.05	1.06	1.07	1.08	1.10	1.08	1.10	1.12
$V_{cb}(10^{-2})$	3.86	3.97	3.96	3.92	3.97	3.97	3.99	4.03	4.01	4.00	4.05	4.06

Table .3. The observable parameters in B^0 and B_s^0 as well as Λ_b decays with $B_{SL} = 10.48\%$.

m_c(GeV)	1.55			1.65			1.75			1.8		
κ_1(GeV^2)	-0.5	-0.6	-0.7	-0.4	-0.5	-0.6	-0.3	-0.4	-0.5	-0.2	-0.3	-0.4
μ(GeV)	2.85	2.91	3.00	2.53	2.62	2.76	2.31	2.43	2.60	2.18	2.30	2.46
$\frac{\tau(B_s^0)}{\tau(B^0)}$	0.904	0.916	0.947	0.915	0.933	0.971	0.927	0.949	0.991	0.927	0.942	0.975
$\frac{\tau(\Lambda_b)}{\tau(B^0)}$	0.704	0.706	0.727	0.723	0.735	0.770	0.747	0.766	0.814	0.755	0.769	0.805
$B_\tau(B^0)$	0.029	0.028	0.028	0.028	0.027	0.027	0.027	0.027	0.027	0.027	0.026	0.026
$\mathcal{R}(B^0)$	0.27	0.27	0.26	0.27	0.26	0.26	0.26	0.25	0.26	0.26	0.25	0.25
$n_c(B^0)$	1.23	1.23	1.24	1.20	1.21	1.22	1.18	1.19	1.21	1.16	1.18	1.20
$B_{SL}(B_s^0)$(%)	10.54	10.58	10.66	10.63	10.69	10.78	10.75	10.82	10.92	10.79	10.86	10.94
$B_\tau(B_s^0)$	0.031	0.030	0.029	0.030	0.029	0.029	0.029	0.029	0.028	0.029	0.029	0.028
$\mathcal{R}(B_s^0)$	0.29	0.28	0.28	0.28	0.28	0.27	0.27	0.27	0.26	0.27	0.27	0.26
$n_c(B_s^0)$	1.22	1.23	1.23	1.19	1.20	1.20	1.16	1.17	1.18	1.14	1.15	1.16
$\frac{B_{SL}(B_s^0)}{B_{SL}(B^0)}$	1.01	1.01	1.02	1.01	1.02	1.03	1.03	1.03	1.04	1.03	1.04	1.04
$B_{SL}(\Lambda_b)$(%)	10.62	10.67	10.77	10.85	10.94	11.09	11.18	11.32	11.51	11.33	11.47	11.65
$B_\tau(\Lambda_b)$	0.035	0.035	0.035	0.035	0.035	0.035	0.035	0.035	0.035	0.036	0.035	0.035
$\mathcal{R}(\Lambda_b)$	0.33	0.33	0.32	0.32	0.32	0.31	0.32	0.31	0.30	0.31	0.31	0.30
$n_c(\Lambda_b)$	1.22	1.22	1.22	1.18	1.18	1.18	1.13	1.13	1.14	1.10	1.11	1.11
$\frac{B_{SL}(\Lambda_b)}{B_{SL}(B^0)}$	1.01	1.02	1.03	1.04	1.04	1.06	1.07	1.08	1.10	1.08	1.09	1.11
$V_{cb}(10^{-2})$	3.75	3.84	3.82	3.85	3.88	3.87	3.96	3.98	3.93	4.01	4.03	4.01

FIGURES

FIG. 1. The lifetime ratios $\tau(\Lambda_b)/\tau(B^0)$ and $\tau(B_s^0)/\tau(B^0)$ as the functions of the running energy scale μ, charm quark mass m_c and κ_1. (1a) and (1c) : $\mu = m_b$, (1b) and (1d) : $m_c = 1.65$GeV. The experimental data are $\tau(\Lambda_b)/\tau(B^0) = 0.79 \pm 0.05$ and $\tau(B_s^0)/\tau(B^0) = 0.94 \pm 0.04$.

FIG. 2. The semileptonic branching ratio B_{SL} as the function of μ, m_c and κ_1. (2a) : $\mu = m_b/2$, (2b) : $m_c = 1.65$GeV and (2c) : $\kappa_1 = -0.6$ GeV^2. The world average is $B_{SL} = 10.48 \pm 0.50\%$.

FIG. 3. B_τ as the function of μ, m_c and κ_1. (3a) : $\mu = m_b/2$, (3b) : $m_c = 1.65$GeV and (3c) : $\kappa_1 = -0.6$ GeV^2. The CLEO data is $B_\tau = 2.6 \pm 0.1\%$.

FIG. 4. R as the function of μ, m_c and κ_1. (4a) : $\mu = m_b$, (4b) : $m_c = 1.65$GeV and (4c) : $\kappa_1 = -0.6$ GeV^2. The CLEO data is $R = 0.25 \pm 0.02$.

FIG. 5. The charm counting n_c as the function of μ, m_c and κ_1. (5a) : $\kappa_1 = -0.6$ GeV^2, (5b) : $\mu = m_b$ and (5c) : $m_c = 1.65$GeV, (d) : $B_{SL} = 10.48\%$. The world average is $n_c = 1.17 \pm 0.04$.

FIG. 6. B_{SL}, B_τ, R and n_c as functions of μ, m_b and $a \equiv m_c/m_b$ in the usual HQET. We have fixed $\lambda_1 = -0.36$ GeV^2. (6a), (6b) and (6d) : $m_b = 4.8$ GeV, (6c) : $\mu = 4.8$ GeV.

FIG. 7. $|V_{cb}|$ as functions of μ, m_c and κ_1. (7a) and (7c) : $\kappa_1 = -0.6$ GeV^2, (7b) : $\mu = m_b/2$, (7d) : correlation between $|V_{cb}|$ and R.

FIG. 8. $|V_{cb}|$ as function of m_b and $a \equiv m_c/m_b$ in usual HQET. With $B_{SL} = 10.48\%$ and $\lambda_1 = -0.36$ GeV^2.

REFERENCES

[1] W.Y. Wang, Y.L. Wu and Y.A. Yan, to be published in Int. J. Mod. Phys. A; hep-ph/9906529, 1999.

[2] Y.L. Wu, Mod. Phys. Lett. **A8**, 819 (1993).

[3] J. Chay, H. Georgi, and B. Grinstein, Phys. Lett. **B247** 399 (1990).

[4] I. Bigi, N.G. Uraltsev, and A. Vainshtein, Phys. Lett. **B293** 430 (1992), Erratum-ibid. **B297** 477 (1993); I. Bigi, M. Shifman, and A. Vainshtein, Phys. Rev. Lett **71** 496 (1993).

[5] B. Blok, L. Koyrakh, M. Shifman, and A.I. Vainshtein, Phys. Rev. **D49** 3356 (1994); Erratum-ibid **D50** 3572 (1994).

[6] A. Manohar and M. Wise, Phys. Rev. **D49** 1310 (1994).

[7] T. Mannel, Nucl. Phys. **B413** 396 (1994).

[8] M. Neubert, Phys. Rev. **D49** 3392 (1994); ibid, 4623.

[9] A. Falk, Z. Ligeti, M. Neubert, and Y. Nir, Phys. Lett. **B326** 145 (1994), and reference therein.

[10] C.C.V. Chang, D. Chang, W.F. Chang, H.N. Li, H.L. Yu, Phys. Rev. **D58** 094019 (1998).

[11] For updated world averages of B hadron lifetimes and lifetime difference, see J. Alcaraz *et al.* (LEP B Lifetime Group), http://wwwcn.cern.ch/~claires/lepblife.html.

[12] H.Y. Cheng and K.C. Yang, Phys. Rev **D59** 014011 (1999).

[13] M. Neubert and C. Sachrajda, Nucl. Phys. **B483** 339 (1997); M. Neubert, hep-ph/9707217.

[14] M. Baek, J. Lee, Chun Liu, and H.S. Song, Phys. Rev. **D57** 4091 (1998), and reference therein.

[15] G. Altarelli, G. Martinelli, S. Petrarca, and F. Rapano, Phys. Lett. **B382** 409 (1996); G. Martinelli, Nucl. Instrum. Methods. Phys. Res. **A384** 241 (1996).

[16] I. Bigi, hep-ph/9612293.

[17] H.Y. Cheng, Phys. Rev. **D56** 2783 (1997).

[18] E. Bagan. P. Ball, M. Braun, and P. Gosdzinsky, Nucl. Phys. **B432** 3 (1994); Phys. Lett. **B342** 362 (1995), Erratum-ibid. **B374** 363 (1996); E. Bagan. P. Ball, B. Fiol, and P. Gosdzinsky, Phys. Lett. **B351** 546 (1995).

[19] M.E. Luke, Phys. Lett. **B252**, 447 (1990).

[20] PDG, Eur. Phys. J. **C3** 1 (1998).

[21] M. Voloshin, Phys. Rev. **D51** 3948 (1995).

[22] A. Czarnecki, M. Jezabek, and J.H. Kuhn, Phys. Lett. **B346** 335 (1995).

[23] M. Luke, M. Savage, and M. Wise, Phys. Lett. **B343** 329 (1995); **B345** 301 (1995).

[24] M. Lu, M. Luke, M. Savage, and B. Smith, Phys. Rev. **D55** 2827 (1997).

[25] CLEO Collaboration, T.E. Coan et.al, Phys. Rev. Lett. **80** 1150 (1998).

[26] I.I. Bigi, M. Shifman, and N. Uraltev, Ann. Rev. Nucl. Part. Sci. **47** 591 (1997) and references therein.

[27] C.Y. Cheung and W.M. Zhang, hep-ph/9810500 and references therein.



$\kappa_1 = -0.6\mathrm{GeV}^2$
$-0.4\mathrm{GeV}^2$
$-0.2\mathrm{GeV}^2$
$\tau(\Lambda_b)/\tau(B^0)$
m_c(GeV)
1.2
1.1
1
0.9
0.8
0.7
1.2
1.3
1.4
1.5
1.6
1.7
1.8
1.9



$\mu=2m_b$
m_b
$m_b/2$
$\tau(\Lambda_b)/\tau(B^0)$
$-\kappa_1(\mathrm{GeV}^2)$
1.2
1.1
1
0.9
0.8
0.7
0
0.1
0.2
0.3
0.4
0.5
0.6
0.7



$\kappa_1 = -0.6\,GeV^2$
$-0.4\,GeV^2$
$-0.2\,GeV^2$
$\tau(B_s)/\tau(B^0)$
m_c(GeV)
1.2
1.15
1.1
1.05
1
0.95
0.9
0.85
1.2
1.3
1.4
1.5
1.6
1.7
1.8
1.9



$\tau(B_s)/\tau(B^0)$
$-\kappa_1(\mathrm{GeV}^2)$
$\mu=2m_b$
m_b
$m_b/2$
1.2
1.15
1.1
1.05
1
0.95
0.9
0.85
0
0.1
0.2
0.3
0.4
0.5
0.6
0.7


κ₁=−0.6GeV²
−0.4GeV²
−0.2GeV²
B_SL
m_c(GeV)
0.08
0.09
0.1
0.11
0.12
0.13
0.14
0.15
1.2
1.3
1.4
1.5
1.6
1.7
1.8
1.9



0.15
0.14
0.13
0.12
0.11
0.1
0.09
0.08
B_{SL}
0
0.1
0.2
0.3
0.4
0.5
0.6
0.7
$-\kappa_1 (GeV^2)$
$\mu = 2m_b$
m_b
$m_b/2$



m_c=1.45GeV
1.65GeV
1.85GeV
B_SL
μ(GeV)
0.08
0.09
0.1
0.11
0.12
0.13
0.14
0.15
2
3
4
5
6
7
8
9
10



B_τ
0.038
0.036
0.034
0.032
0.03
0.028
0.026
0.024
1.2
1.3
1.4
1.5
1.6
1.7
1.8
1.9
m_c(GeV)
$\kappa_1 = -0.6\text{GeV}^2$
-0.4GeV^2
-0.2GeV^2



B_τ
0.024
0.026
0.028
0.03
0.032
0.034
0.036
0.038
0
0.1
0.2
0.3
0.4
0.5
0.6
0.7
$-\kappa_1(GeV^2)$
$\mu=2m_b$
m_b
$m_b/2$



0.038
0.036
0.034
0.032
0.03
0.028
0.026
0.024
B_τ
2
3
4
5
6
7
8
9
10
μ(GeV)
m_c=1.45GeV
1.65GeV
1.85GeV


0.38
0.36
0.34
0.32
0.3
0.28
0.26
0.24
0.22
0.2
0.18
R
1.2
1.3
1.4
1.5
1.6
1.7
1.8
1.9
m_c(GeV)
$\kappa_1 = -0.6\text{GeV}^2$
-0.4GeV^2
-0.2GeV^2



R
0.18
0.2
0.22
0.24
0.26
0.28
0.3
0.32
0.34
0.36
0.38
0
0.1
0.2
0.3
0.4
0.5
0.6
0.7
$-\kappa_1(\mathrm{GeV}^2)$
$\mu=2m_b$
m_b
$m_b/2$


R
0.38
0.36
0.34
0.32
0.3
0.28
0.26
0.24
0.22
0.2
0.18
2
3
4
5
6
7
8
9
10
μ(GeV)
m_c=1.45GeV
1.65GeV
1.85GeV



$\mu = m_b/2$
m_b
$2m_b$
n_c
1.35
1.3
1.25
1.2
1.15
1.1
1.05
1.2
1.3
1.4
1.5
1.6
1.7
1.8
1.9
m_c(GeV)



n_c
1.35
1.3
1.25
1.2
1.15
1.1
1.05
0
0.1
0.2
0.3
0.4
0.5
0.6
0.7
$-\kappa_1(GeV^2)$
m_c=1.45GeV
1.65GeV
1.85GeV



$\kappa_1=-0.6\mathrm{GeV}^2$
$-0.4\mathrm{GeV}^2$
$-0.2\mathrm{GeV}^2$
n_c
1.35
1.3
1.25
1.2
1.15
1.1
1.05
2
3
4
5
6
7
8
9
10
μ(GeV)


n_c
m_c(GeV)
1.05
1.1
1.15
1.2
1.25
1.3
1.35
1.2
1.3
1.4
1.5
1.6
1.7
1.8
1.9
κ_1=−0.2GeV²
−0.4GeV²
−0.6GeV²


B_SL
μ(GeV)
0.15
0.14
0.13
0.12
0.11
0.1
0.09
0.08
2
3
4
5
6
7
8
9
10
a=0.34
0.30
0.26



B_τ
0.04
0.035
0.03
0.025
0.02
0.015
2
3
4
5
6
7
8
9
10
μ(GeV)
a=0.26
0.30
0.34


R
a
0.34
0.32
0.3
0.28
0.26
0.24
0.22
0.2
0.18
0.16
0.14
0.26
0.27
0.28
0.29
0.3
0.31
0.32
0.33
0.34
m_b=5.0GeV
4.8GeV
4.6GeV


$\mu=m_b/2$
m_b
$2m_b$
n_c
a
1.05
1.1
1.15
1.2
1.25
1.3
1.35
0.26
0.27
0.28
0.29
0.3
0.31
0.32
0.33
0.34



0.048
0.046
0.044
0.042
0.04
0.038
0.036
0.034
0.032
|V_cb|
1.2
1.3
1.4
1.5
1.6
1.7
1.8
1.9
m_c(GeV)
B_SL=0.0998
0.1048
0.1098


0.048
0.046
0.044
0.042
0.04
0.038
0.036
0.034
0.032
$|V_{cb}|$
$m_c = 1.85$GeV
1.65GeV
1.45GeV
0
0.1
0.2
0.3
0.4
0.5
0.6
0.7
$-\kappa_1(\mathrm{GeV}^2)$



0.048
0.046
0.044
0.042
0.04
0.038
0.036
0.034
0.032
$|V_{cb}|$
2
2.5
3
3.5
4
4.5
5
μ(GeV)
m_c=1.45GeV
1.65GeV
1.85GeV



$|V_{cb}|$
R
0.048
0.046
0.044
0.042
0.04
0.038
0.036
0.034
0.032
0.23
0.24
0.25
0.26
0.27
0.28
0.29
0.3
0.31
0.32
$\kappa_1 = -0.6\,\mathrm{GeV}^2$
$-0.4\,\mathrm{GeV}^2$
$-0.2\,\mathrm{GeV}^2$



0.048
0.046
0.044
0.042
0.04
0.038
0.036
0.034
0.032
$|V_{cb}|$
0.26
0.27
0.28
0.29
0.3
0.31
0.32
a
m_b=4.7GeV
4.8GeV
4.9GeV



a=0.32
0.29
0.26
|V_cb|
0.048
0.046
0.044
0.042
0.04
0.038
0.036
0.034
0.032
4.6
4.65
4.7
4.75
4.8
4.85
4.9
4.95
5
m_b